

Albania -- Nursing Students Global Health Professionals Survey (GHPS) FACT SHEET

The Albania Nursing Students GHPS includes data on prevalence of cigarette smoking and other tobacco use, knowledge and attitudes about tobacco use, exposure to second-hand smoke, desire for smoking cessation, and training received during nursing school regarding patient counseling on smoking cessation techniques. This information could be very useful in monitoring and developing Albania's comprehensive tobacco control program.

The Albania Nursing Students GHPS was a school-based survey of all 3rd year nursing students attending the 5 nursing schools in Albania. The school response rate was 100% and the student response rate was 68.2%. A total of 356 students participated in the Albania Nursing Students GHPS.

Prevalence

62.6% ever smoked cigarettes (male = 80.6%; female = 57.4%).
41.5% currently smoked cigarettes (male = 57.5%; female = 36.4%).
7.9% ever used other tobacco products (male = 22.7%; female = 3.4%).
1.5% currently used other tobacco products (male = 2.1%; female = 1.3%).
29.1% of ever smokers smoked on school premises/property during past year.
34.4% of ever smokers smoked in school buildings during past year.
20.2% of ever other tobacco users used other tobacco products on school premises/property during past year.
17.6% of ever other tobacco users used other tobacco products in school buildings during past year.

Factors Influencing Tobacco Use

79.2% were exposed to second-hand smoke at home, during past week.
93.8% were exposed to second-hand smoke in public places, during past week.
24.1% thought school has ban on smoking in school buildings and clinics.
29.5% thought school enforces the ban on smoking in school buildings and clinics.
11.6% thought school has ban on smoking in indoor public or common areas.
26.9% thought school enforces the ban on smoking in indoor public or common areas.
94.3% thought smoking should be banned in all enclosed public places.
96.9% thought tobacco sales to adolescents should be banned.
66.9% thought there should be a complete ban on advertising of tobacco products.
87.0% of current smokers wanted to stop smoking.
54.3% of current smokers ever received help/advice to stop smoking cigarettes.
89.2% were taught about dangers of smoking during nursing training.

Role Model and Cessation Training

82.3% thought health professionals serve as role models for their patients.
22.6% received formal training in smoking cessation approaches during nursing school.
96.7% thought health professionals should get specific training on cessation techniques.

Highlights

Out of 100 students...

- **42 currently smoked cigarettes and 2 used other tobacco products.**
- **37 of 42 smokers wanted to quit.**
- **94 were exposed to smoke in public places.**
- **94 supported a ban on smoking in all enclosed public places.**
- **23 received training to provide patients with cessation approaches.**
- **97 thought health professionals should get specific training on cessation techniques to use with patients.**

For additional information, please contact:

Roland Shuperka Shuperka2004@yahoo.com
Arta Lena atlandus@yahoo.com